

Quantization of generalized Virasoro-like algebras ¹

Guang'ai Song*, Yucai Su[†] and Yuezhu Wu[‡]

* College of Mathematics and Information Science, Shandong Institute of Business
and Technology, Yantai, Shandong 264005, China

[†] Department of Mathematics, University of Science and Technology of China,
Hefei 230026, China

[‡] Department of Mathematics, Qufu Normal University, Qufu, Shandong 273165, China

E-mail: gasong@ccec.edu.cn

Abstract. In a recent paper by the authors, Lie bialgebras structures of generalized Virasoro-like type were considered. In this paper, the explicit formula of the quantization of generalized Virasoro-like algebras is presented.

Key words: quantization, Lie bialgebras, Drinfel'd twist, generalized Virasoro-like algebras

§1. Introduction

In Hopf algebras or quantum groups theory, there are two standard methods to yield new bialgebras from old ones, one is twisting the product by a 2-cocycle but keeping the coproduct unchanged, another is twisting the coproduct by a Drinfel'd twist element but keeping the product unchanged. Constructing quantization of Lie bialgebras is an important method to produce new quantum groups (cf. [V], [PO], etc). In the paper [M1] (cf. [M2, M3]), a class of infinite dimensional Lie bialgebras containing Virasoro algebras was presented. This type Lie bialgebras was classified in [NT], and the quantization of this type Lie algebras was determined in [G]. In the paper [SS], Lie bialgebras structures of generalized Witt type were classified. The quantization of this type algebras was considered in [HW]. In the paper [WSS], the structures of Lie bialgebras of generalized Virasoro-like algebras were determined. In the preset paper, we will consider the quantization of this type Lie algebras.

§2. Preliminaries

§2.1 Generalized Virasoro-like Lie bialgebras. Let $\mathbb{C}$ denote the complex field, and let Γ be any nondegenerate additive subgroup of $\mathbb{C}^2$ (namely Γ contains a $\mathbb{C}$ -basis of $\mathbb{C}^2$).

¹Supported by NSF grant Y2006A17 of Shandong Province, NSF J06P52 of Shandong Provincial Education Department, China, NSF grant 10171064 of China, "One hundred Talents Program" from University of Science and Technology of China

Definition 2.1 The Lie algebras $\mathcal{L}(\Gamma)$ with basis $\{L_\alpha, \partial_1, \partial_2 | \alpha \in \Gamma \setminus \{0\}\}$ and bracket:

$$[L_\alpha, L_\beta] = (\alpha_1\beta_2 - \alpha_2\beta_1)L_{\alpha+\beta}, [\partial_i, L_\alpha] = \alpha_i L_{\alpha_i}, \text{ for } \alpha, \beta \in \Gamma \setminus \{0\}, i = 1, 2.$$

is called a generalized Virasoro-like algebras.

Remark: We use the convention that if an undefined notation appears in an expression, we always treat it as zero; for instance, $L_\alpha = 0$, if $\alpha = 0$. In particular, when $\Gamma = \mathbb{Z}^2$ the derived subalgebras $[\mathcal{L}(\mathbb{Z}^2), \mathcal{L}(\mathbb{Z}^2)] = \text{span}\{L_\alpha | \alpha \in \mathbb{Z}^2 \setminus \{0\}\}$ is the Virasoro-like algebras (cf. [LT, MJ, ZZ]). The Lie algebras $\mathcal{L}(\Gamma)$ is closely related to the Lie algebras of Block type (cf. [DZ, X, Z]) and the Lie algebras of Cartan type S (cf. [SX, X1, Z2]).

The following theorem is the main result in [WSS].

Theorem 2.2 Every Lie bialgebra structure on the Lie algebras $\mathcal{L}(\Gamma)$ is a coboundary triangular Lie bialgebra, namely, it is given by r -matrix $r - r^{21}$ with $r = T \otimes L_\alpha, T \in \text{span}\{\alpha_1, \alpha_2\}, L_\alpha \in \mathcal{L}(\Gamma)$.

§2.2 Drinfel'd twisting. Let A be a unital R -algebra (where R is a ring), for any element $x \in A, a \in R$, we set

$$x_a^{\langle n \rangle} = (x + a)(x + a + 1) \cdots (x + a + n - 1), \quad (2.1)$$

$$x_a^{[n]} = (x + a)(x + a - 1) \cdots (x + a - n + 1), \quad (2.2)$$

where $n \in \mathbb{Z}$, and we denote $x^{\langle n \rangle} = x_0^{\langle n \rangle}, x^{[n]} = x_0^{[n]}$.

The following lemma belongs to [G] and [GZ].

Lemma 2.3 Let $\mathbb{F}$ be a field with $\text{char } \mathbb{F} = 0$, and x be any element of a unital $\mathbb{F}$ -algebras A , for $a, d \in \mathbb{F}$, and $m, n, r \in \mathbb{Z}$, one has:

$$x_a^{\langle m+n \rangle} = x_a^{\langle m \rangle} x_{m+a}^{\langle n \rangle}, \quad (2.3)$$

$$x_a^{[m+n]} = x_a^{[m]} x_{a-m}^{[n]}, \quad (2.4)$$

$$x_a^{[m]} = x_{a-m+1}^{\langle m \rangle}, \quad (2.5)$$

$$\sum_{m+n=r} \frac{(-1)^n}{m!n!} x_a^{[m]} x_d^{\langle n \rangle} = \binom{a-d}{r} = \frac{(a-d)(a-d+1) \cdots (a-d-r+1)}{r!}, \quad (2.6)$$

$$\sum_{m+n=r} \frac{(-1)^m}{m!n!} x_a^{[m]} x_{d-m}^{[n]} = \binom{a-d+r-1}{r} = \frac{(a-d)(a-d+1) \cdots (a-d+r-1)}{r!}. \quad (2.7)$$

The following definition belongs to [D].

Definition 2.4 An element $\mathcal{F} \in \mathcal{H} \otimes \mathcal{H}$ is called Drinfel'd twisting element, if it is invertible such that

$$(\mathcal{F} \otimes 1)(\Delta_0 \otimes Id)(\mathcal{F}) = (1 \otimes \mathcal{F})(1 \otimes \Delta_0)(\mathcal{F}) \quad (2.8)$$

$$(\epsilon_0 \otimes Id)(\mathcal{F}) = 1 \otimes 1 = (Id \otimes \epsilon_0)(\mathcal{F}). \quad (2.9)$$

The following theorem is well known (cf.[D]), it can be found in any book of hopf algebras.

Theorem 2.5 Let $(H, \mu, \tau, \Delta_0, \epsilon_0, S_0)$ be a hopf algebra over commutative ring, $\mathcal{F}$ be a Drinfel'd element of $H \otimes H$, then

(1) $u = \mu(Id \otimes S_0)(\mathcal{F})$ is a invertible element of $H \otimes H$ with $u^{-1} = \mu(S_0 \otimes Id)(\mathcal{F})$.

(2) The algebras $(H, \mu, \tau, \Delta, \epsilon, S)$ is a new hopf algebra if we keep the counit undeformed and define $\Delta : H \rightarrow H \otimes H$, $S : H \rightarrow H$ by:

$$\Delta(h) = \mathcal{F}\Delta_0(h)\mathcal{F}^{-1}, \quad S(h) = uS_0(h)u^{-1}.$$

Let $(\mathcal{U}(\mathcal{L}(\Gamma)), \mu, \tau, \Delta_0, \epsilon_0, S_0)$ be the standard hopf algebra, i.e.

$$\Delta_0(L_\alpha) = L_\alpha \otimes 1 + 1 \otimes L_\alpha, \quad \Delta_0(\partial_i) = \partial_i \otimes 1 + 1 \otimes \partial_i,$$

$$S(L_\alpha) = -L_\alpha, \quad S(\partial_i) = -\partial_i, \quad \epsilon_0(L_\alpha) = 0, \quad \epsilon_0(\partial_i) = 0,$$

for $\alpha \in \mathbb{F} \setminus \{0\}, i = 1, 2$. The main result of this paper is the following theorem.

Theorem 2.6 Let $\mathcal{L}(\Gamma)$ be the generalized Virasoro-like algebra over $\mathbb{F}$ with $\text{char } \mathbb{F} = 0$, choose $T = a_1\partial_1 + a_2\partial_2 \in \text{span}\{\partial_1, \partial_2\}$ and $L_\alpha \in \mathcal{L}(\Gamma)$ with $[T, L_\alpha] = L_\alpha$, then there exists a noncommutative and noncocommutative hopf algebra structure $(\mathcal{U}(\mathcal{L}(\Gamma))[[t]], \mu, \tau, \Delta, S, \epsilon)$ on $\mathcal{U}(\mathcal{L}(\Gamma))[[t]]$ over $\mathbb{F}[[t]]$, such that $\mathcal{U}(\mathcal{L}(\Gamma))[[t]]/t\mathcal{U}(\mathcal{L}(\Gamma))[[t]] = \mathcal{U}(\mathcal{L}(\Gamma))$, which preserves the product and the counit of $\mathcal{U}(\mathcal{L}(\Gamma))[[t]]$, but the coproduct and antipode are defined by

$$\Delta(L_\beta) = L_\beta \otimes (1 - L_\alpha t)^b + \sum_{i=0}^{\infty} (-1)^i T^i \otimes (1 - L_\alpha t)^{-i} L_{\beta+i\alpha} c_i t^i, \quad (2.10)$$

$$\Delta(\partial_i) = \partial_j \otimes 1 + 1 \otimes \partial_j + \alpha_j T^{(1)} \otimes (1 - L_\alpha t)^{-1} L_\alpha t, \quad j = 1, 2. \quad (2.11)$$

$$S(L_\beta) = -(1 - L_\alpha t)^{-b} \sum_{i=0}^{\infty} L_{\beta+i\alpha} c_i T_1^{(i)} t^i, \quad (2.12)$$

$$S(\partial_i) = \alpha_j T (1 - L_\alpha t)^{-1} (L_\alpha t - L_\alpha^2 t^2) - \partial_j \quad (2.13)$$

where $b = a_1\beta_1 + a_2 + \beta_2$, $c_i = \frac{(\alpha_1\beta_2 - \alpha_2\beta_1)^i}{i!}$, $c_0 = 1$, $j = 1, 2$.

§3. Proof of the main result

We shall divide the proof of the Theorem 2.6 into several lemmas.

Lemma 3.1 Let $T, L_\alpha \in \mathcal{L}(\Gamma)$ with

$$T = a_1 \partial_1 + a_2 \partial_2, \quad [T, L_\alpha] = [a_1 \partial_1 + a_2 \partial_2, L_\alpha] = L_\alpha,$$

for some $a_1, a_2 \in \mathbb{F}$ and any $\alpha = (\alpha_1, \alpha_2) \in \Gamma$. For any $\beta = (\beta_1, \beta_2) \in \Gamma$, denote $b = a_1 \beta_1 + a_2 \beta_2$. The following equations hold in $\mathcal{U}(\mathcal{L}(\Gamma))$ for $a \in \mathbb{F}$, $m, k \in \mathbb{Z}_+$:

$$L_\beta T_a^{[m]} = T_{a-b}^{[m]} \beta_\beta, \quad (3.1)$$

$$L_\beta T_a^{\langle m \rangle} = T_{a-b}^{\langle m \rangle} L_\beta, \quad (3.2)$$

$$L_\alpha^k T_a^{[m]} = T_{a-k}^{[m]} L_\alpha^k, \quad (3.3)$$

$$L_\alpha^k T_a^{\langle m \rangle} = T_{a-k}^{\langle m \rangle} L_\alpha^k, \quad (3.4)$$

$$\partial_j^k T_a^{[m]} = T_a^{[m]} \partial_j^k, j = 1, 2, \quad (3.5)$$

$$\partial_j^k T_a^{\langle m \rangle} = T_a^{\langle m \rangle} \partial_j^k, j = 1, 2, \quad (3.6)$$

$$L_\beta L_\gamma^m = \sum_{i=0}^m (-1)^i \binom{m}{i} (\gamma_1 \beta_2 - \gamma_2 \beta_1)^i L_\gamma^{m-i} L_{\beta+i\gamma}, \quad (3.7)$$

$$\partial_j L_\gamma^m = m \gamma_j L_\gamma^m + L_\gamma^m \partial_j, j = 1, 2. \quad (3.8)$$

Proof. Since

$$[T, L_\beta] = [a_1 \partial_1 + a_2 \partial_2, L_\beta] = (a_1 \beta_1 + a_2 \beta_2) L_\beta = b L_\beta = T L_\beta - L_\beta T,$$

we have $L_\beta T = (T - b) L_\beta$, it easy to see that (2.1) is true for $m = 1$. Suppose that (3.1) is true for m , then for $m + 1$ we have

$$L_\beta T_a^{[m+1]} = L_\beta T_a^{[m]} (T + a - m) = T_{a-b}^{[m]} L_\beta (T + a - m) = T_{a-b}^{[m]} (T + a - b - m) L_\beta = T_{a-b}^{[m+1]}.$$

By induction on m , (3.1) holds. Similarly, we can obtain (3.2)–(3.6). For (3.7), we have:

$$\begin{aligned} L_\beta L_\gamma^m &= \sum_{i=0}^m (-1)^i \binom{m}{i} L_\gamma^{m-i} (\text{ad} L_\gamma)^i (L_\beta) \\ &= \sum_{i=0}^m (-1)^i \binom{m}{i} L_\gamma^{m-i} L_{\beta+i\gamma} (\gamma_1 \beta_2 - \gamma_2 \beta_1)^i. \end{aligned}$$

The proof of formula (3.8) is similar to that of (3.7), using the following fact:

$$\text{ad}(L_\gamma)^i (\partial_j) = \begin{cases} -\gamma_j L_\gamma, & \text{if } i = 1, \\ 0, & \text{if } i > 1. \end{cases} \quad \square$$

Now for $a \in \mathbb{F}$, set

$$\begin{aligned}\mathcal{F}_a &= \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_a^{[i]} \otimes L_\alpha^i t^i, \\ F_a &= \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_a^{(i)} \otimes L_\alpha t^i, \\ u_a &= \mu \cdot (S_0 \otimes Id)(F_a), v_a = \mu \cdot (Id \otimes S_0)(\mathcal{F}_a).\end{aligned}$$

Write $\mathcal{F} = \mathcal{F}_0$, $F = F_0$, $u = u_0$, $v = v_0$. Since $S_0(T_a^{(i)}) = (-1)^i T_{-a}^{[i]}$, $S_0(L_\alpha^i) = (-1)^i L_\alpha$, we have:

$$\begin{aligned}u_a &= \mu(S_0 \otimes Id) \left(\sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_\alpha^i t^i \right) = \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_{-a}^{[i]} L_\alpha^i t^i, \\ v_a &= \mu(Id \otimes S_0) \left(\sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_a^{[i]} \otimes L_\alpha^i t^i \right) = \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{[i]} L_\alpha^i t^i.\end{aligned}$$

Lemma 3.2. For $a, d \in \mathbb{F}$, one has

$$\mathcal{F}_a F_d = 1 \otimes (1 - L_\alpha t)^{(a-d)}, v_a u_d = (1 - L_\alpha t)^{-(a+d)}.$$

Therefore the elements $\mathcal{F}_a, F_a, u_a, v_a$ are invertible elements with $\mathcal{F}_a^{-1} = F_a$, $u_a^{-1} = v_{-a}$.

Proof. Using the formula (2.6) we have:

$$\begin{aligned}\mathcal{F}_a F_d &= \left(\sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_a^{[i]} \otimes L_\alpha^i t^i \right) \cdot \left(\sum_{j=0}^{\infty} \frac{1}{j!} T_d^{(j)} \otimes L_\alpha^j t^j \right) \\ &= \sum_{i,j=0}^{\infty} \frac{(-1)^i}{i!j!} T_a^{[i]} T_d^{(j)} \otimes L_\alpha^i L_\alpha^j t^{i+j} = \sum_{m=0}^{\infty} \left(\sum_{i+j=m} \frac{(-1)^j}{i!j!} T_a^{[i]} T_d^{(j)} \right) \otimes L_\alpha^m t^m \\ &= \sum_{m=0}^{\infty} (-1)^m \binom{a-d}{m} L_\alpha^m t^m = 1 \otimes (1 - L_\alpha t)^{a-d}, \\ v_a u_d &= \left(\sum_{i=0}^{\infty} \frac{1}{i!} T_a^{[i]} L_\alpha^i t^i \right) \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} T_{-d}^{[j]} L_\alpha^j t^j = \sum_{i,j=0}^{\infty} \frac{1}{i!j!} T_a^{[i]} L_\alpha^i t^i \frac{(-1)^j}{j!} T_{-d}^{[j]} L_\alpha^j t^j \\ &= \sum_{m=0}^{\infty} \sum_{i+j=m} \frac{(-1)^j}{i!j!} T_a^{[i]} T_{-d}^{[j]} L_\alpha^{i+j} t^{i+j} \quad (\text{from (3.3)}) \\ &= \sum_{m=0}^{\infty} \binom{a+d+m-1}{m} L_\alpha^m t^m \quad (\text{from (2.7)}) \\ &= (1 - L_\alpha t)^{-(a+d)}.\end{aligned}$$

□

Lemma 3.3. For any nonnegative integer m , and any $a \in \mathbb{F}$ we have

$$\Delta_0 T^{[m]} = \sum_{i=0}^m \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]}.$$

In particular, we have $\Delta_0 T^{[m]} = \sum_{i=0}^m \binom{m}{i} T^{[i]} \otimes T^{[m-i]}$.

Proof. Since $\Delta_0(T) = T \otimes 1 + 1 \otimes T$, it is easy to see that the result is true for $m = 1$. Suppose it is true for m , then for $m + 1$, we have:

$$\begin{aligned}
\Delta_0(T^{[m+1]}) &= \Delta_0(T^{[m]})\Delta_0(T - m) \\
&= \left(\sum_{i=0}^m \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]}\right)((T - a - m) \otimes 1 + 1 \otimes (T + a - m) + m(1 \otimes 1)) \\
&= \left(\sum_{i=1}^{m-1} \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]}\right)((T - a - m) \otimes 1 + 1 \otimes (T + a - m)) \\
&\quad + m\left(\sum_{i=0}^m \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]}\right) + (1 \otimes T_a^{[m+1]} + T_{-a}^{[m+1]} \otimes 1) \\
&\quad + (T - a - m) \otimes T_a^{[m]} + T_{-a}^{[m]} \otimes (T + a - m) \\
&= 1 \otimes T_a^{[m+1]} + T_{-a}^{[m+1]} \otimes 1 + m\left(\sum_{i=1}^{m-1} \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]}\right) \\
&\quad + (T - a) \otimes T_a^{[m]} + T_{-a}^{[m]} \otimes (T + a) + \sum_{i=1}^{m-1} \binom{m}{i} T_{-a}^{[i+1]} \otimes T_a^{[m-i]} \\
&\quad + \sum_{i=1}^{m-1} (i - m) \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]} + \sum_{i=1}^{m-1} \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i+1]} \\
&\quad + \sum_{i=1}^{m-1} (-i) \binom{m}{i} T_{-a}^{[i]} \otimes T_a^{[m-i]} \\
&= 1 \otimes T_a^{[m+1]} + T_{-a}^{[m+1]} \otimes 1 + \sum_{i=1}^m \left(\binom{m}{i-1} + \binom{m}{i} \right) T_{-a}^{[i]} \otimes T_a^{[m+1-i]} \\
&= \sum_{i=0}^{m+1} \binom{m+1}{i} T_{-a}^{[i]} \otimes T_a^{[m+1-i]}.
\end{aligned}$$

From induction the result hold on arbitrary m . □

Lemma 3.3. $\mathcal{F} = \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T^{[i]} \otimes L_{\alpha}^i t^i$ is a Drinfel'd twist element of $\mathcal{U}(\mathcal{L}(\Gamma))[[t]]$, i.e.

$$(\mathcal{F} \otimes 1)(\Delta_0 \otimes Id)(\mathcal{F}) = (1 \otimes \mathcal{F})(1 \otimes \Delta_0)(\mathcal{F}),$$

$$(\epsilon_0 \otimes Id)(\mathcal{F}) = 1 \otimes 1 = (Id \otimes \epsilon_0)(\mathcal{F}).$$

Proof. The second equation holds obviously, we just need to prove the first one. Since

$$\begin{aligned}
(\mathcal{F} \otimes 1)(\Delta_0 \otimes Id)(\mathcal{F}) &= \left(\sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T^{[i]} \otimes L_{\alpha}^i t^i \otimes 1 \right) (\Delta_0 \otimes Id) \left(\sum_{j=0}^{\infty} \frac{(-1)^j}{j!} T^{[j]} \otimes L_{\alpha}^j t^j \right) \\
&= \left(\sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T^{[i]} \otimes L_{\alpha}^i t^i \otimes 1 \right) \times \\
&\quad \times \left(\sum_{j=0}^{\infty} \frac{(-1)^j}{j!} \sum_{k=0}^j \binom{j}{k} T_{-i}^{[k]} \otimes T_i^{[j-k]} \otimes L_{\alpha}^j t^j \right) \\
&= \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j}}{i!j!} t^{i+j} \sum_{k=0}^j \binom{j}{k} T^{[i]} T_{-i}^{[k]} \otimes L_{\alpha}^i T_i^{[j-k]} \otimes L_{\alpha}^j \\
&= \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j}}{i!j!} t^{i+j} \sum_{k=0}^j \binom{j}{k} T^{[i+k]} \otimes T^{[j-k]} L_{\alpha}^i \otimes L_{\alpha}^j,
\end{aligned}$$

on the other hand,

$$\begin{aligned}
(1 \otimes \mathcal{F})(Id \otimes \Delta_0)(\mathcal{F}) &= \left(\sum_{r=0}^{\infty} \frac{(-1)^r}{r!} t^r 1 \otimes T^{[r]} \otimes L_{\alpha}^r \right) \times \\
&\quad \times \left(\sum_{s=0}^{\infty} \frac{(-1)^s}{s!} t^s T^{[s]} \otimes \sum_{q=0}^s \binom{s}{q} L_{\alpha}^q \otimes L_{\alpha}^{s-q} \right) \\
&= \sum_{r,s=0}^{\infty} \frac{(-1)^{r+s}}{r!s!} t^{r+s} \sum_{q=0}^s \binom{s}{q} T^{[s]} \otimes T^{[r]} L_{\alpha}^q \otimes L_{\alpha}^{r+s-q},
\end{aligned}$$

it sufficient to show for a fixed m that:

$$\sum_{i+j=m} \frac{1}{i!j!} t^{i+j} \sum_{k=0}^j \binom{j}{k} T^{[i+k]} \otimes T^{[j-k]} L_{\alpha}^i \otimes L_{\alpha}^j = \sum_{r+s=m} \frac{1}{r!s!} t^{r+s} \sum_{q=0}^s \binom{s}{q} T^{[s]} \otimes T^{[r]} L_{\alpha}^q \otimes L_{\alpha}^{r+s-q}.$$

Now fix r , $0 \leq i \leq s$, and set $i = q$, $i + k = s$, then we have

$$j - k = j - (s - i) = i + j - s = m - s = r.$$

We see that the coefficients of $T^{[s]} \otimes T^{[r]} L_{\alpha}^q \otimes L_{\alpha}^{m-q}$ in both sides are equal. So the result holds. $\square$

Lemma 3.4. For $a \in \mathbb{F}$, $\beta \in \Gamma$, we have

$$(L_{\beta} \otimes 1)F_a = F_{a-b}(L_{\beta} \otimes 1), \quad (3.9)$$

$$(1 \otimes L_{\beta})F_a = \sum_{l=0}^{\infty} (-1)^l F_{a+l} (T_a^{(l)} \otimes c_l L_{\beta+l\alpha} t^l), \quad (3.10)$$

$$L_\beta u_a = u_{a+b} \sum_{l=0}^{\infty} L_{\beta+l\alpha} c_l T_{-a+l}^{(l)} t^l, \quad (3.11)$$

$$(\partial_j \otimes 1)F_a = F_a(\partial_j \otimes 1), \quad (3.12)$$

$$(1 \otimes \partial_j)F_a = F_{a+1}T_a^{(1)} \otimes \alpha_j L_\alpha t + F_a(1 \otimes \partial_j), \quad (3.13)$$

$$\partial_j u_a = -\alpha_j T_{-a}^{[1]} u_{a+1} L_\alpha t + u_a \partial_j, \quad (3.14)$$

$$L_\alpha u_a = u_{a+1} L_\alpha, \quad (3.15)$$

$$v T_{-a}^{[1]} = T_{-a}^{[1]} v_{-a} - T_a^{[1]} v_a L_\alpha t. \quad (3.16)$$

where $c_l = \frac{1}{l!}(\alpha_1 \beta_2 - \alpha_2 \beta_1)^l$, $j = 1, 2$.

Proof. From (3.2) we have:

$$\begin{aligned} (L_\beta \otimes 1)F_a &= (L_\beta \otimes 1) \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_\alpha^i t^i \\ &= \sum_{i=0}^{\infty} \frac{1}{i!} L_\beta T_a^{(i)} \otimes L_\alpha^i t^i \\ &= \sum_{i=0}^{\infty} \frac{1}{i!} T_{a-b}^{(i)} L_\beta \otimes L_\alpha^i t^i = F_{a-b}(L_\beta \otimes 1), \end{aligned}$$

where $b = \alpha_1 \beta_2 + \alpha_2 \beta_1$. This proves (3.9). For (3.10), using (3.7) we have:

$$\begin{aligned} (1 \otimes L_\beta)F_a &= (1 \otimes L_\beta) \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_\alpha^i t^i = \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_\beta L_\alpha^i t^i \\ &= \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes \sum_{l=0}^i (-1)^l \binom{i}{l} L_\alpha^{i-l} L_{\beta+l\alpha} (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l t^i \\ &= \sum_{i=0}^{\infty} \left(\sum_{l=0}^i (-1)^l \frac{1}{(i-l)! l!} T_a^{(i)} \otimes L_\alpha^{i-l} L_{\beta+l\alpha} (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l \right) t^i \\ &= \sum_{i=0}^{\infty} \sum_{l=0}^i (-1)^l \frac{1}{i! l!} T_a^{(i+l)} \otimes L_\alpha^i L_{\beta+l\alpha} (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l t^{i+l} \\ &= \sum_{l=0}^{\infty} (-1)^l \sum_{i=0}^{\infty} \left(\frac{1}{i!} T_{a+l}^{(i)} \otimes L_\alpha^i t^i \right) \frac{1}{l!} T_a^{(l)} \otimes L_{\beta+l\alpha} t^l (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l \\ &= \sum_{l=0}^{\infty} (-1)^l F_{a+l} (T_a^{(l)} \otimes L_{\beta+l\alpha} c_l t^l). \end{aligned}$$

So we have (3.10). Now we prove (3.11):

$$\begin{aligned} L_\beta u_a &= L_\beta \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a}^{[r]} L_\alpha^r t^r = \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} L_\beta T_{-a}^{[r]} L_\alpha^r t^r = \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a-b}^{[r]} L_\beta L_\alpha^r t^r \\ &= \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a-b}^{[r]} \sum_{l=0}^r (-1)^l \binom{r}{l} L_\alpha^{r-l} (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l L_{\beta+l\alpha} t^r \\ &= \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a-b}^{[r]} \sum_{l=0}^r (-1)^l \frac{r!}{(r-l)! l!} L_\alpha^{r-l} (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l L_{\beta+l\alpha} t^r \end{aligned}$$

$$\begin{aligned}
&= \sum_{r,l=0}^{\infty} \frac{(-1)^r}{r! l!} T_{-a-b}^{[r+l]} L_{\alpha}^r (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l L_{\beta+l\alpha} t^{r+l} \\
&= \sum_{r,l=0}^{\infty} \frac{(-1)^r}{r! l!} T_{-a-b}^{[r]} T_{-a-b-r}^{[l]} L_{\alpha}^r (\alpha_1 \beta_2 - \alpha_2 \beta_1)^l L_{\beta+l\alpha} t^{r+l} \\
&= \sum_{l=0}^{\infty} \sum_{r=0}^{\infty} \left(\frac{(-1)^r}{r!} T_{-a-b}^{[r]} L_{\alpha}^r t^r \right) T_{-a-b}^{[l]} L_{\beta+l\alpha} \frac{(\alpha_1 \beta_2 - \alpha_2 \beta_1)^l}{l!} t^l \\
&= u_{a+b} \sum_{l=0}^{\infty} T_{-a-b}^{[l]} L_{\beta+l\alpha} c_l t^l = u_{a+b} \sum_{l=0}^{\infty} L_{\beta+l\alpha} T_{-a+l}^{[l]} c_l t^l = u_{a+b} \sum_{l=0}^{\infty} L_{\beta+l\alpha} T_{1-a}^{(l)} c_l t^l.
\end{aligned}$$

This proves (3.11). For (3.12) we have:

$$\begin{aligned}
(\partial_j \otimes 1)F_a &= (\partial_j \otimes 1) \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_{\alpha}^i t^i = \sum_{i=0}^{\infty} \frac{1}{i!} \partial_j T_a^{(i)} \otimes L_{\alpha}^i t^i \\
&= \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \partial_j \otimes L_{\alpha}^i t^i = \left(\sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_{\alpha}^i t^i \right) (\partial_j \otimes 1) = F_a (\partial_j \otimes 1).
\end{aligned}$$

Using (2.3) and (3.8) we have:

$$\begin{aligned}
(1 \otimes \partial_j)F_a &= (1 \otimes \partial_j) \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_{\alpha}^i t^i = \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes \partial_j L_{\alpha}^i t^i \\
&= \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes (i\alpha_j L_{\alpha}^i + L_{\alpha}^i \partial_j) t^i \\
&= \sum_{i=0}^{\infty} \frac{1}{(i-1)!} T_a^{(i)} \otimes \alpha_j L_{\alpha}^i t^i + \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{(i)} \otimes L_{\alpha}^i \partial_j t^i \\
&= \sum_{i=0}^{\infty} \frac{1}{(i-1)!} T_a^{(1)} T_{a+1}^{(i-1)} \otimes \alpha_j L_{\alpha}^i t^i + F_a (1 \otimes \partial_j) \\
&= F_{a+1} (T_a^{(1)} \otimes \alpha_j L_{\alpha} t) + F_a (1 \otimes \partial_j).
\end{aligned}$$

So (3.13) holds. For (3.14) we have:

$$\begin{aligned}
\partial_j u_a &= \partial_j \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a}^{[r]} L_{\alpha}^r t^r = \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a}^{[r]} \partial_j L_{\alpha}^r t^r \\
&= \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a}^{[r]} (r\alpha_j L_{\alpha}^r + L_{\alpha}^r \partial_j) t^r \\
&= \sum_{r=0}^{\infty} \alpha_j T_{-a}^{[1]} \frac{(-1)^r}{(r-1)!} T_{-a-1}^{[r-1]} L_{\alpha}^r t^r + \sum_{r=0}^{\infty} \frac{(-1)^r}{r!} T_{-a}^{[r]} L_{\alpha}^r t^r \partial_j \\
&= -\alpha_j T_{-a}^{[1]} u_{a+1} L_{\alpha} t + u_a \partial_j.
\end{aligned}$$

Now we prove (3.15):

$$L_{\alpha} u_a = L_{\alpha} \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_{-a}^{[i]} L_{\alpha}^i t^i = \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} T_{-a-1}^{[i]} L_{\alpha}^{i+1} t^i = u_{a+1} L_{\alpha}.$$

For the last equation we have:

$$\begin{aligned}
vT_{-a}^{[1]} &= \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{[i]} L_{\alpha}^i t^i T_{-a}^{[1]} = \sum_{i=0}^{\infty} \frac{1}{i!} T_a^{[i]} (T - a - i) L_{\alpha}^i t^i \\
&= T_{-a}^{[1]} v_a - \sum_{i=0}^{\infty} \frac{1}{(i-1)!} (T + a) T_{a-1}^{[i-1]} L_{\alpha}^i t^i \\
&= T_{-a}^{[1]} v_a - T_a^{[1]} v_{a-1} L_{\alpha} t.
\end{aligned}$$

This complete the proof of the lemma. $\square$

Proof of Theorem 2.6. For arbitrary elements $L_{\beta}, \partial_j \in \mathcal{L}(\Gamma), j = 1, 2$, we have:

$$\begin{aligned}
\Delta(L_{\beta}) &= \mathcal{F} \Delta_0(L_{\beta}) \mathcal{F}^{-1} = \mathcal{F}(L_{\beta} \otimes 1) \mathcal{F}^{-1} + \mathcal{F}(1 \otimes L_{\beta}) \mathcal{F}^{-1} \\
&= \mathcal{F}(L_{\beta} \otimes 1) F + \mathcal{F}(1 \otimes L_{\beta}) F \\
&= \mathcal{F} F_{-b}(L_{\beta} \otimes 1) + \mathcal{F} \sum_{l=0}^{\infty} (-1)^l F_l(T^{(l)} \otimes L_{\beta+l\alpha} c_l t^l) \\
&= (1 \otimes (1 - L_{\alpha} t)^b)(L_{\beta} \otimes 1) \\
&\quad + \sum_{l=0}^{\infty} (-1)^l (1 \otimes (1 - L_{\alpha} t)^{-l}) \otimes (T^{(l)} \otimes L_{\beta+l\alpha} c_l t^l) \\
&= L_{\beta} \otimes (1 - L_{\alpha} t)^b + \sum_{l=0}^{\infty} (-1)^l T^{(l)} \otimes (1 - L_{\beta} t)^{-l} L_{\beta+l\alpha} c_l t^l,
\end{aligned}$$

$$\begin{aligned}
\Delta(\partial_j) &= \mathcal{F} \Delta(\partial_j) \mathcal{F}^{-1} = \mathcal{F}(\partial_j \otimes 1 + 1 \otimes \partial_j) F \\
&= \mathcal{F}(\partial_j \otimes 1) F + \mathcal{F}(1 \otimes \partial_j) F \\
&= \mathcal{F} F(\partial_j \otimes 1) + \mathcal{F}(F_1(T^{(1)} \otimes \alpha_j L_{\alpha} t) + F(1 \otimes \partial_j)) \\
&= \partial_j \otimes 1 + 1 \otimes \partial_j + 1 \otimes (1 - L_{\alpha} t)^{-1} (T^{(1)} \otimes \alpha_j L_{\alpha} t) \\
&= \partial_j \otimes 1 + 1 \otimes \partial_j + \alpha_j T^{(1)} \otimes (1 - L_{\alpha} t)^{-1} L_{\alpha} t, j = 1, 2,
\end{aligned}$$

$$\begin{aligned}
S(L_{\beta}) &= u^{-1} S_0(L_{\beta}) u = -v L_{\beta} u = -v u_b \left(\sum_{l=0}^{\infty} c_l L_{\beta+l\alpha} T_1^{(l)} t^l \right) \\
&= -(1 - L_{\alpha} t)^{-b} \left(\sum_{l=0}^{\infty} c_l L_{\beta+l\alpha} T_1^{(l)} t^l \right),
\end{aligned}$$

$$\begin{aligned}
S(\partial_j) &= u^{-1} S_0(\partial) u = -v \partial_j u \\
&= -v(-\alpha_j T^{[1]} u_1 L_{\alpha} t + u \partial_j) \\
&= \alpha_j (T v - T v L_{\alpha} t) u_1 L_{\alpha} t - \partial_j \\
&= \alpha_j T v u_1 L_{\alpha} t - \partial_j T v u_2 L_{\alpha}^2 t^2 - \partial_j \\
&= \alpha_j T (1 - L_{\alpha} t)^{-1} L_{\alpha} t - \alpha_j T (1 - L_{\alpha} t)^{-1} L_{\alpha}^2 t^2 - \partial_j \\
&= \alpha_j T (1 - L_{\alpha} t)^{-1} (L_{\alpha} t - L_{\alpha}^2 t^2) - \partial_j.
\end{aligned}$$

$\square$

References

[D1] V. Drinfel'd, Constant quasiclassical solutions of the Yang-Baxter quantum equation, *Soviet*

- Math. Dokl.* **28**(3) (1983), 667–671.
- [D2] V. Drinfel'd Quantum groups, Proceedings ICM (Berkeley 1986), Providence, Amer Math Soc, 1987, 789–820.
- [F] R. Farnsteiner, Derivations and central extensions of finitely generalized Lie algebras, *J. Algebra* **118** (1988), 33–45.
- [GZ] A. Giaquinto, J. Zhang, Bialgebra action, twists and universal deformation formulas, *J. Pure Appl. Algebra* **128**(2) (1998), 133–151.
- [G] C. Grunspan, Quantizations of the Witt algebra and of simple Lie algebras in characteristic p , *J. Algebra* **280** (2004), 145–161.
- [HW] N. Hu, X. Wang, Quantizations of generalized-Witt algebra and of Jacobson-Witt algebra in modular case, arXiv:math.QA/0602281.
- [M1] W. Michaelis, A class of infinite-dimensional Lie bialgebras containing the Virasoro algebras, *Adv. Math.* **107** (1994), 365–392.
- [M2] W. Michaelis, Lie coalgebras, *Adv. Math.* **38** (1980), 1–54.
- [M3] W. Michaelis, The dual Poincare-Birkhoff-Witt theorem, *Adv. Math.* **57** (1985), 93–162.
- [NT] S.-H. Ng, Earl J. Taft, Classification of the Lie bialgebra structures on the Witt and Virasoro algebras, *J. P. App. algebra.* **151** (2000), 67–88.
- [N] W.D. Nichols, The structure of the dual Lie coalgebra of the Witt algebra, *J. P. APP. Alg.* **68** (1990), 395–364W
- [P] D. Passman, New Simple Infinite Dimensional Lie Algebras, *J. Algebra* **206** (1998), 682–692.
- [SS] G. Song, Y. Su, Lie bialgebras of generalized Witt type, *Science in China, Series A–Mathematics* **49**(4) (2006), 533–544.
- [T] Earl J. Taft, Witt and Virasoro algebras as Lie bialgebras, *J. P. App. algebra.* **87** (1993), 301–312.
- [WSS] Y. Wu, G. song, Y. Su, Lie bialgebras of generalized Virasoro-like type, *Acta Mathematica Sinica, English Series* **22**(6) (2006), 1915–1922.